

L15 ANSWER 54 OF 101 CA COPYRIGHT 2002 ACS  
 AN 105:95850 CA  
 TI Complementation of class II A alleles in the immune response to  
 (GluLysTyr) polymers  
 AU Lai, C. H.; Babu, U. M.; Matsunaga, K.; Nagy, Z. A.; Klein, J.; Turchin,  
 H. A.; Maurer, P. H.  
 CS Jefferson Med. Coll., Thomas Jefferson Univ., Philadelphia, PA, 19107, USA  
 SO Exp. Clin. Immunogenet. (1986 ), 3(1), 38-48  
 CODEN: ECIME4  
 DT Journal  
 LA English  
 AB The proliferative T cell responses to poly(GluLysTyr) (GLT) and  
 poly(GluLysPhe) (GLPhe) are restricted by the E.alpha.E.beta. class II MHC  
 mol. (E) in most responder mouse strains. Some nonresponder strains that  
 carry responder E.beta., but cannot express cell surface E mols., can  
 complement with other nonresponder strains that provide the missing  
 E.alpha. chain needed for the expression of E mols. and for responsiveness  
 to GLT and GLPhe. Here, another type of complementation is described  
 between 2 E-nonexpressor haplotypes, H-2f and H-2s, which result in  
 E-nonexpressor F1 hybrids, which are responders to GLT. The restriction  
 element involved in this response is an Af/Ashybrid mol. The data support  
 the hypothesis that conformational determinants resulting from the free  
 assocn. of .alpha. and .beta. chains in heterozygotes can increase the  
 immune potential of the individual.

Applicants: Alexander Gad and Dora Lis  
 Serial No.: 09/816,989  
 Filed: March 23, 2001  
 Exhibit 28